

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 10 *Swine Day (1968-2014)*

Article 465

1991

On-farm feed uniformity survey

C R. Stark

J A. Hansen

Robert D. Goodband

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

 Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Stark, C R.; Hansen, J A.; Goodband, Robert D.; and Behnke, Keith C. (1991) "On-farm feed uniformity survey," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.6305>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1991 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



On-farm feed uniformity survey

Abstract

An on-farm feed manufacturing survey was conducted in conjunction with a mixer uniformity analysis. The survey collected information on producer size and type of operation, mixer type, manufacturing practices, and feed production. The survey represents 43 participants from across the U.S. Results indicate that 42% of the participants had mixer uniformity coefficients of variation of less than 10%, 47% were between 10-20%. and 11 % had CV's greater than 20%.; Swine Day, Manhattan, KS, November 21, 1991

Keywords

Swine day, 1991; Kansas Agricultural Experiment Station contribution; no. 92-193-S; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 641; Swine; On-farm feed manufacturing; Feed uniformity; Mixers

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Authors

C R. Stark, J A. Hansen, Robert D. Goodband, and Keith C. Behnke

K

S

U

ON-FARM FEED UNIFORMITY SURVEY

*C. R. Stark^{1,2}, K. C. Behnke²,
R. D. Goodband, and J. A. Hansen*

Summary

An on-farm feed manufacturing survey was conducted in conjunction with a mixer uniformity analysis. The survey collected information on producer size and type of operation, mixer type, manufacturing practices, and feed production. The survey represents 43 participants from across the U.S. Results indicate that 42% of the participants had mixer uniformity coefficients of variation of less than 10%, 47% were between 10-20%, and 11% had CV's greater than 20%.

(Key Words: On-Farm Feed Manufacturing, Feed Uniformity, Mixers.)

Procedures

Mixer analysis kits were sent out upon request to producers to test their mixing efficiency. The producer had to obtain 10 samples at evenly spaced intervals as the feed was removed from the mixer. In addition, each producer had to complete a survey form. Information was collected on producer size and type of operation, mixer type, manufacturing practices, and feed production.

The samples were analyzed for uniformity of salt distribution by the Quantab^R Ion strip test. Coefficient of variation (CV) was calculated for each producer. When using the Quantab^R Ion test, CV's less than 10% indicated a uniformly mixed diet (a CV of 10% takes into account sampling errors and analytical errors).

Results and Discussion

Results indicate that a majority of on-farm feed manufacturers may not be producing a uniformly mixed feed (Table 1) with desired CV's of less than 10%. Non-uniformity becomes more critical when medicated feed additives are included in the feed. The survey indicated that production size had no significant effect on uniformity results. Results did not favor any particular mixer type.

Feed uniformity can be improved by proper ingredient sequencing. A proper ingredient sequence would be: protein source, premix, flush after the premix (protein source or ground grain), then grain. Grinding to a particle size of less than 800 microns also will help improve feed uniformity. Periodic inspec-

¹To obtain a mixer analysis kit send your name and address to:

Charles Stark
Shellenberger Hall
Kansas State University
Manhattan, KS 66506

²Department of Grain Science and Industry.

tion and repair of worn equipment will help prevent poor feed uniformity. Finally, mixers should be tested annually to determine if the feed manufacturing protocol is sufficient or if

it should be changed. However, the first step is to develop a feed manufacturing protocol that assures a uniform feed.

Table 1. Results of Mixer Uniformity Analysis for Each Mixer Type

Mixer Type	Coefficient of Variation ^a		
	< 10 %	10-20 %	20+ %
Vertical Portable	13	13	3
Vertical Stationary	1	1	1
Horizontal Paddle	2	2	--
Horizontal Double Ribbon	2	--	1
Drum	--	2	--
Other	--	2	--
Total	18	20	5

^aA coefficient of variation less than 10% indicates a uniformly mixed diet.



Lyell Nelson, Dr. Don Kropf, and John Wolf evaluate carcasses in the KSU Meats Lab,